

## CENTER FOR COLON & RECTAL HEALTH, INC.

Richard Goldstein MD, Anne-Marie Marcoux MD, David Schaffzin MD  
215-741-4910

### MOVIPREP DAY BEFORE

**TWO WEEKS BEFORE** your procedure **stop** taking all **herbal vitamins, vitamin E, and fish oil.**

**IF YOU ARE TAKING A BLOOD THINNER** under a doctor's order, you must check with your doctor before stopping that medication.

**IF YOU ARE DIABETIC**, speak to your doctor about your medication dosage on the day you are prepping and also on the day of your procedure.

**ONE WEEK BEFORE** your procedure **stop** taking any of the following medications and any medication containing **aspirin or ibuprofen:**

Advil	Clinoril	Heparin	Norgesic
Aleve	Coumadin	Ibuprofen	Nuprin
Alka Seltzer	Daypro	Indocin	Percodan
Anacin	Dipridamole	Lodine	Persantine
Anaprox	Disalcid	Lovenox	Plavix
Arthropan	Ecotrin	Meclomen	Relafen
Ascriptin	Empirin	Mobic	Ticlid
Aspirin	Excedrin	Motrin	Toradol
Baby Aspirin	Feldene	Naprosyn	Voltaren
Bufferin	Fiorinal	Naproxen	

**\*\*\* TYLENOL IS SAFE TO TAKE \*\*\***

**FOR FIVE DAYS BEFORE** your procedure try to avoid eating vegetables with skin such as corn, peas, beans, etc. (If you do eat them, you may still have your procedure.)

\*\* You may want to purchase diaper rash cream or vaseline to apply while prepping. \*\*

1. **At least three days before your procedure** fill your prescription for Moviprep.
2. **On the day before your procedure** you may only have clear liquids the entire day. This includes: water, soda of any kind, and clear juices such as apple, grape, cranberry. You **may not have citrus juices** such as orange juice or grapefruit juice. You may have jello (but **no red jello**). You may have clear broth or bouillon, and tea and coffee with sugar or sugar substitutes. You may have a small splash of milk in tea and coffee in AM only.
3. **At 8:00 AM** the day before your procedure empty (1) Pouch A and (1) Pouch B into the disposable container. Add lukewarm drinking water to the top of the fill line. Shake the container. Refrigerate until ready to use. The solution must be used within 24 hours.
4. **At 3:00 PM** shake the solution prior to drinking. The Moviprep container is divided by 4 marks. Every 15 minutes drink the solution down to the next mark (approximately 8 oz.) until the full liter is completed. **As soon as you finish the first liter**, mix the second dose following the above directions and refrigerate. Before completing the next step, drink an additional 16 oz. of clear liquids to ensure hydration.
5. **At 6:00 PM** begin drinking the second container of Moviprep as shown in step 4. If you are unable to finish the solution or have any problems with the prep, please contact our office. You may continue to drink clear liquids up until midnight. The more clear liquids you drink, the better the prep will work. Then have **nothing after midnight** (including no gum, no hard candy or mints, etc.).
6. **On the day of your procedure** take only your necessary morning medications (blood pressure, heart, asthma, etc.) with a small sip of water **when you wake up**. All other medications may be taken when you get home.
7. Please wear loose-fitting clothing.
8. Someone **must drive you home** after your procedure. You may not take a taxi.
9. If there is a change in your insurance or health history (heart attack, breathing problems, etc.), you must notify us immediately. If you need to reschedule your appointment, please give us as much notice as possible.