

**CENTER FOR COLON AND RECTAL HEALTH, INC.**  
Richard Goldstein MD, Anne-Marie Marcoux MD, David Schaffzin MD  
215-741-4910

**SUPREP SPLIT DAY PROCEDURE**

**TWO WEEKS BEFORE** your procedure **stop** taking all **herbal vitamins, vitamin E, and fish oil.**

**IF YOU ARE TAKING A BLOOD THINNER** under a doctor's order, you must check with your doctor before stopping that medication.

**IF YOU ARE DIABETIC,** speak to your doctor about your medication dosage on the day you are prepping and also on the day of your procedure.

**ONE WEEK BEFORE** your procedure **stop** taking any of the following medications and any medication containing **aspirin or ibuprofen:**

Advil	Clinoril	Heparin	Norgesic
Aleve	Coumadin	Ibuprofen	Nuprin
Alka Seltzer	Daypro	Indocin	Percodan
Anacin	Dipridamol	Lodine	Persantine
Anaprox	Disalcid	Lovenox	Plavix
Arthropan	Ecotrin	Meclomen	Relafen
Ascriptin	Empirin	Mobic	Ticlid
Aspirin	Excedrin	Motrin	Toradol
Baby Aspirin	Feldene	Naprosyn	Voltaren
Bufferin	Fiorinal	Naproxen	

**\*\*\* TYLENOL IS SAFE TO TAKE \*\*\***

**FOR FIVE DAYS BEFORE** your procedure try to avoid eating vegetables with skin such as corn, peas, beans, etc. (If you do eat them, you may still have your procedure.)

**\*\* You may want to purchase diaper rash cream or vaseline to apply while prepping. \*\***

- 1. At least two days before your procedure** fill your prescription for Suprep.
- 2. On the day before your procedure** you may have breakfast and a light lunch.  
**After lunch** you may have **only clear liquids** which include: water, soda of any kind, and clear juices such as apple, grape, cranberry. **You may not have citrus juices** such as orange juice or grapefruit juice. You may have jello ( **but no red jello**). You may have bouillon, clear broth, and tea and coffee with sugar or sugar substitutes.  
After lunch you may **not** have milk in tea and coffee.
- 3. At 6:00 PM** pour one 6 oz. bottle of Suprep into the container. Fill the rest of the container to the 16 oz. line with cool water, seven-up or gingerale (diet is OK) and drink all of the liquid in the container. After finishing the Suprep you **must drink** two (2) more 16 oz. containers of liquid **over the next hour**. You may drink more clear liquids the rest of the evening.
- 4. At 6:00 AM on the day of your procedure** repeat step # 3 with the second bottle of Suprep. You may have clear liquids **until 9:00 AM** and then **nothing after 9:00 AM** (including no gum, no hard candy, or mints, etc.).
- 5. At 8:30 AM on the day of your procedure** take only your necessary morning medications (heart, blood pressure, asthma, etc.). All other medications may be taken when you get home.
- 6.** Please wear loose-fitting clothing.
- 7.** Someone **must drive you home** after your procedure. You may not take a taxi.
- 8.** If there is a change in your insurance or health history (heart attack, breathing problems, etc.) you must notify us immediately. If you need to reschedule your appointment, please give us as much notice as possible.